



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07K 7/06, A01N 63/02, C12N 1/20 // (C12N 1/20, C12R 1:125)	A1	(11) International Publication Number: WO 98/50422 (43) International Publication Date: 12 November 1998 (12.11.98)
(21) International Application Number: PCT/US98/09471 (22) International Filing Date: 8 May 1998 (08.05.98) (30) Priority Data: 08/853,753 9 May 1997 (09.05.97) US (71) Applicant (for all designated States except US): AGRAQUEST, INC. [US/US]; Suite 4, 1105 Kennedy Place, Davis, CA 95616 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): HEINS, Sherry, Darlene [US/US]; 3110 Oyster Bay, Davis, CA 95616 (US). MANKER, Denise, Carol [US/US]; 3037 Prado Lane, Davis, CA 95616 (US). JIMENEZ, Desmond, Rito [US/US]; 9 Pendergast Street, Woodland, CA 95695 (US). McCOY, Randy, Jay [US/US]; 3212 Breton Avenue, Davis, CA 95616 (US). ORJALA, Jimmy, Ensio [US/US]; 2905 Lollard Drive, Davis, CA 95616 (US). (74) Agents: PARK, Freddie, K. et al.; Morrison & Foerster LLP, 755 Page Mill Road, Palo Alto, CA 94304-1018 (US).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, GW, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: A NOVEL STRAIN OF <i>BACILLUS</i> FOR CONTROLLING PLANT DISEASES AND CORN ROOTWORM		
(57) Abstract <p>The present invention relates to a novel antibiotic-producing and metabolite-producing <i>Bacillus subtilis</i> strain that exhibits insecticidal, antifungal and antibacterial activity. The supernatant of this novel strain contains effective insecticidal, antifungal and antibacterial agents. Also included in the invention is a solvent extractable, small molecular weight (< 10,000 daltons) corn rootworm-active metabolite produced in the supernatant. Also included in the invention are methods of protecting or treating plants from fungal and bacterial infections and corn rootworm infestations comprising the step of applying to the plant an effective amount of the antibiotic/metabolite-producing novel <i>Bacillus subtilis</i> strain, the antibiotic/metabolite produced by the novel <i>Bacillus subtilis</i> strain or a combination thereof, optionally further comprising another antibiotic-producing bacterial strain and/or a chemical pesticide. The invention also includes methods of preventing or treating fungal and bacterial infections using whole broth cultures or supernatants obtained from cultures of the novel <i>Bacillus subtilis</i> strain alone or in combination with chemical pesticides and/or other biocontrol agents. The invention also includes novel antifungal and antibacterial compounds designated agrastatins and a novel combination comprising an A-type iturin, a plipastatin, a surfactin and an agrastatin. Methods of treating or protecting plants from fungal and bacterial infections and corn rootworm infestations comprising administering the novel agrastatins and the novel combination comprising an A-type iturin, a plipastatin, a surfactin and an agrastatin are provided.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

5

A NOVEL STRAIN OF *BACILLUS* FOR CONTROLLING PLANT DISEASES AND CORN ROOTWORM

10

Field of the Invention

The present invention is in the field of biopesticides. More particularly, this invention relates to the finding that a novel strain of *Bacillus subtilis*, AQ713, can inhibit a broad range of fungal and bacterial plant diseases and also have activity against corn rootworm. The invention also relates to fungicidal, bactericidal, and insecticidal compositions comprising this novel *Bacillus* strain and the antibiotics and metabolites produced by this strain either alone, or in combination with other chemical and biological pesticides.

15

Cross Reference to Related Applications

20

This application is a continuation-in-part of Serial No. 08/853,753, filed May 9, 1997.

Background of the Invention

25

For a number of years, it has been known that various microorganisms exhibit biological activity so as to be useful to control plant diseases. Although progress has been made in the field of identifying and developing biological pesticides for controlling various plant diseases of agronomic and horticultural importance, most of the pesticides in use are still synthetic compounds. Many of these chemical fungicides are classified as carcinogens by the EPA, are toxic to wildlife and other non-target species. In addition, pathogens may develop

30

resistance to chemical pesticides (see, e.g., Schwinn *et al.*, p. 244, ADVANCES IN PLANT PATHOLOGY: *PHYTOPHTHORA INFESTANS*, THE CAUSE OF LATE BLIGHT OF POTATO (Academic Press, San Diego 1991).

Every year 250-300 million dollars of chemical pesticides are used to control corn rootworm infestations. Many of these chemical pesticides are toxic to humans, wildlife and other nontarget species. Also some have been found in the ground water. New chemical insecticides cost \$100 million to develop.

Biological control offers an attractive alternative to synthetic chemical fungicides. Biopesticides (living organisms and the naturally produced compounds produced by these organisms) can be safer, more biodegradable, and less expensive to develop.

Screening programs have identified certain *Bacillus* spp. (*Bacillus* spp. includes *B. subtilis*, *B. cereus*, *B. mycoides*, *B. thuringiensis*) strains that exhibit antifungal activity. (See e.g. Stabb *et al.* (1990) *Applied Environ. Microbiol.* 60: 4404-4412). These strains have been shown to produce zwittermicin-A and or kanosamine (Milner *et al.* (1996) *Appl. Environ. Microb.* 62: 3061-3066), two antibiotic agents that are effective against the soil borne disease damping off, caused by *Phytophthora medicaginis*, *P. nicotianae*, *P. aphanidermatum* or *Sclerotinia minor* (See Stabb *et al.*, *supra*). Zwittermicin-A is a water soluble, acid stable linear aminopolyol molecule (see, He *et al.*, (1994) *Tetra. Lett.* 35 (16) 2499-2502.

U.S. Patent No. 5,049,379 to Handelsman *et al.* describes how zwittermicin-A produces damping off in alfalfa and soybeans. When the seed was coated with *B. cereus* ATCC 53522, the pathogenic activity of root rot fungus is inhibited. Similarly application of spore-based formulations of certain *B. cereus* strains to soybean seeds or the soil surrounding the seeds has been shown to improve soybean yield at field sites. (See, Osburne *et al.* (1995) *Am. Phytopathol. Soc.* 79(6): 551-556). Methods of applying biopesticides are well known in the art and include, for example, wettable powders, dry flowables, microencapsulation of effective agents, liquid or solid formulations of antibiotic fractions from

suitable cultures. (See e.g., U.S. Patent No. 5,061,495 to Rossall or U.S. Patent No. 5,049,379 to Handelsman).

Smith *et al.* (1993) *Plant Disease* 77(2) 139-142 report that the activity of the soil-borne fungus, *Pythium aphanidermatum*, that causes cottony cucumber leak can be suppressed using zwittermicin-producing *B. cereus* strain UW85. Leifert *et al.* (1995) *J. Appl. Bacteriol.* 78: 97-108 report that the production of anti-*Botrytis* and anti-*Alternaria* antibiotics by two *Bacillus* strains, *B. subtilis* CL27 and *B. pumilis* CL 45. The whole broth and cell-free filtrates were active against *Botrytis* and *Alternaria* in in vitro tests and were active against *Botrytis* in in vivo small plant tests on *Astilbe*. Leifert *et al.* (1997) U.S. Patent No. 5,597,565 disclose *B. subtilis*, *B. pumilis*, and *B. polymyxa* that are particularly effective at inhibiting post harvest disease causing fungi. They also disclose the presence of antibiotics produced in the cell-free culture filtrate and their activity at different pH values, but they do not identify these compounds.

Rossall (1994) U.S. Patent No. 5,344,647 discloses *Bacillus subtilis* strains with broad anti-fungal activity. Sholberg *et al.* (1995) *Can. J. Microbiol.* 41: 247-252, Swinburne *et al.* (1975) *Trans. Brit. Mycol. Soc.* 65: 211-217, Singh and Deverall (1984) *Trans. Br. Mycol. Soc.* 83: 487-490, and Ferreira, *et al.* (1991) *Phytopathology* 81: 283-287. Baker *et al.* (1983) *Phytopathology* 73: 1148-1152 disclose the use of *Bacillus* spp. and *Bacillus subtilis* as biocontrol agents of fungal plant pathogens. Baker *et al.* (1983) *Phytopathology* 73: 1148-1152 also report on an antifungal *Bacillus subtilis* for use on plant pathogens. Pusey *et al.* (1988) *Plant Dis.* 72: 622-626, Pusey and Robins (U.S. Patent No. 5,047,239), and McKeen *et al.* (1986) *Phytopathology* 76: 136-139 disclose control of post harvest fruit rot using *B. subtilis*. McKeen *et al.*, *supra*, have shown that antibiotics similar to the low molecular weight iturin cyclic polypeptides contribute to this fungicidal activity of *B. subtilis*.

Liu *et al.* (1995) U.S. Patent No. 5,403,583 disclose a *Bacillus megaterium*, ATCC 55000 and a method to control the fungal plant pathogen, *Rhizoctonia solani*. Islam and Nandi (1985) *Journal of Plant Diseases and*

Protection 92(3): 241-246 disclose a *Bacillus megaterium* with antagonism to *Drechslera oryzae*, the causal agent of rice brown spot. The same authors, Islam and Nandi (1985) *Journal of Plant Diseases and Protection* 92(3) 233-240 also disclose in-vitro antagonism of *B. megaterium* against *Drechslera oryzae*,
5 *Alternaria alternata* and *Fusarium roseum*. They discuss three components in the culture filtrate. The most active antibiotic was highly soluble in water and methanol with a UV peak at 255 nm and a shoulder at 260 nm, which proved to be a polyoxin-like lipopeptide. Cook ((1987) *Proceedings Beltwide Cotton Production - Mechanization Research Conference*, Cotton Council, Memphis, p.
10 43-45) discloses the use of a suspension of *Bacillus megaterium* to reduce the number of cotton plants killed by *Phymatotrichum omnivorum*, a cause of cotton root rot.

Antibiotic production of *B. megaterium* has been recorded by Berdy (CRC Handbook of Antibiotic Compounds, Vols. I-XIV, (CRC Press, Inc. Boca Raton,
15 FL 1980-87) who reports production of low-mammalian toxic peptide antibiotics such as ansamitocin-PDM-O, bacimethrin, megacin, pentapeptide, homopeptides.


Bacilli are known to produce antifungal and antibacterial secondary metabolites (Korzybski *et al.* (1978)). University of Wisconsin and Cornell researchers have identified a novel fungicidal compound, zwittermicin A,
20 produced by *Bacillus sp.* (He *et al.* (1994) *Tetra. Lett.* 35(16):2499-2502). A second fungicidal metabolite produced by the same strain was recently identified as the known amino-sugar, kanosamine (Milner *et al.* (1996) *Appl. Environ. Microb.* 62:3061-3065).

Another group of previously described *Bacillus* metabolites are the cyclic
25 lipopeptides of the iturin class, some of which are potent fungicidal agents. These agents consist of a cyclic octapeptide with seven α -amino acids and one β -amino acid with an aliphatic side chain. There are several groups of iturins that differ in order and content of the amino acid sequence. These are shown in Table 1 below. Generally, a suite of related molecules is produced with differences in the length
30 and branching of the aliphatic amino acid residue. When tested against

Saccharomyces cerevesiae, mycosubtilin was found to be the most active agent (LC50 = 10 µg/mL) followed by iturin-A and bacillomycin L (both having an LC50 = 30 µg/mL) (Beeson *et al.* (1979) *J. Antibiotics* 32(8):828-833). The mode of action of these cyclic lipopeptides has been reported to be due to interaction with fungal membranes creating transmembrane channels that permit release of vital ions (Latoud *et al.* (1986) *Biochem. Biophys. Acta* 856:526-535). Iturin-C is inactive against fungi including *Penicillium chrysogenum* (Peypoux *et al.* (1978) *Tetrahedron* 34:1147-1152).

Table 1
Structures of the iturin family of antibiotics

Antibiotic	L-Asz(X1)	X4	X5	X6	X7
Iturin A	L-Asn	L-Gln	L-Pro	D-Asn	L-Ser
Iturin C	L-Asp	L-Gln	L-Pro	D-Asn	L-Ser
Bacillo-mycin D	L-Asn	L-Pro	L-Glu	D-Ser	L-Thr
Bacillo-mycin L	L-Asp	L-Ser	L-Gln	D-Ser	L-Thr
Bacillo-mycin F	L-Asn	L-Gln	L-Pro	D-Asn	L-Thr
Myco-subtilin	L-Asn	L-Gln	L-Pro	D-Ser	L-Asn



A research group at the USDA has investigated the structure/activity relationship of the iturins by synthesizing a number of analogs differing in the amino acid chain length. The researchers reported that the activity of the iturins increased with the length of the fatty acid side chain and the terminal branching in the order iso>normal>anteiso (Bland *et al.* (1995) *Proc. Plant Growth Regulation Soc. Am.* 22nd: 105-107). They also state that the "amounts of iturins obtained from natural production are inadequate to be commercially viable" based on their work with a number of iturin producing strains of *Bacillus*.

Another groups of cyclic lipopeptides isolated from *B. cereus* are the plipastatins. These compounds are a family of acylated decapeptides, the structures of which are shown in Figure 1 (Nishikiori *et al.* (1986) *J. Antibiotics* 39(6):755-761). These compounds were originally isolated as inhibitors of porcine pancreatic phospholipase A₂ (Umezawa *et al.* (1986) *J. Antibiotics* 39(6):737-744), but were later found to inhibit some plant pathogenic fungi including *Botrytis*, *Pyricularia* and *Alternaria* (Yamada *et al.* (1990) *Nippon Noyaku Gakkaishi* 15(1):95-96). Yamada also reported a synergistic effect observed between iturin A and the plipastatins, both produced by the same *B. subtilis* strain.

Work has been carried out on fermentation improvements to increase production of the iturins in both liquid (Phae and Shoda (1991) *J. Ferment. Bioeng.* 71:118-121); Ohno *et al.* (1993) *J. Ferment. Bioeng.* 75:463-465) and solid state fermentations (Ohno *et al.* (1992) *Biotech. Lett.* 14(9):817-822; Ohno *et al.* (1995) *J. Ferment. Bioeng.* 5:517-519). There is a report of synergy between the closely related surfactins, that are themselves inactive, and the iturins produced by the same *B. subtilis* strain (Hiraoka *et al.* (1992) *J. Gen. Appl. Microbiol.* 38:635-640). The nucleotide sequence for the gene that co-regulates biosynthesis of iturin A and surfactin has been published (Huang *et al.* (1993) *J. Ferment. Bioeng.* 76(6):445-450). Field work on iturin-producing strains has concentrated on soil treatment for control of *Rhizoctonia* (Asaka and Shoda

(1996) *Appl. Environ. Microbiol.* 62:4081-4085) and foliar field applications of iturins have not been reported.

Another cyclic lipopeptide compound produced by *B. subtilis* is surfactin, which possesses an exceptional surfactant activity (Kaninuma *et al.* (1969) *Agric. Biol. Chem.* 33:973-976). Surfactin contains a C14 or C15 β -hydroxy fatty acid linked by a lactone ring to a heptapeptide moiety with a LLDLLDL sequence (Arima *et al.* (1968) *Biochem. Biophys. Res. Commun.* 31:488-494. Sandrin *et al.* ((1990) *Biotechnol. Appl. Biochem.* 12:370-375) found *B. subtilis* strains that produced both surfactin and iturin A, the bacillomycins F and L and mycosubtilin.

The novel microorganism AQ713 discovered by the inventors, previously thought to be a strain of *Bacillus megaterium* and now identified as a strain of *Bacillus subtilis*, produces A iturins, plipastatins and surfactins. Production of this combination of lipopeptides by a microorganism has not been previously reported. In addition, the inventors have discovered that AQ713 also produces a newly described group of compounds designated as "agrastatins." The combination of all three of the above known compounds with the novel agrastatins is also novel.

One commonly used biopesticide is the gram positive bacterium *Bacillus thuringiensis*. Pesticidal *B. thuringiensis* strains are known to produce crystal proteins during sporulation, which are specifically toxic to certain orders and species of insects and nematodes (See, *e.g.*, U.S. Patents Nos. 4,999,192 and 5,208,017). Proteinaceous endotoxins produced by *B. thuringiensis* also act as insecticidal agents against corn rootworm and other beetles (*e.g.*, U.S. Patent No. 5,187,091; Johnson, T.J. *et al.* (1993), *J. Economic Entomology* 86:330-333). *B. thuringiensis* endotoxins have been shown to be effective as purified crystals, washed cell pellets, and expressed proteins. Warren *et al.* (WO 96/10083), discloses non-endotoxin proteins produced during the vegetative stage of *Bacillus cereus* and *B. thuringiensis*. These vegetative proteins, called Vip1 and Vip2 have potent activity against corn rootworm (northern and western) (Estruch *et al.*

(1997), *Nature Biotechnology* 15:137-141 and Mullins *et al.* (1997), *Appl Environ. Microbiol.* 63, (in press).

One *B. thuringiensis* thermostable metabolites, termed beta-exotoxin has also been shown to have pesticidal properties. Burgjeron and Biache (1979),
5 *Entomophaga* 11:279-284 report a beta exotoxin that is active against Colorado potato beetle (*Leptinotarsa decemlineata*). In addition, the known *B. thuringiensis* beta-exotoxins exhibits non-specific pesticidal activity, killing not only nematodes, but also flies, armyworms, mites, and corn rootworms. Sigma exotoxin has a structure similar to beta-exotoxin, and is active against Colorado
10 potato beetle (Argauer *et al.* (1991) *J. Entomol. Sci.* 26:206-213). Alpha-exotoxin is toxic against larvae of *Musca domestica* (Cluthy (1980) *FEMS Microbiol. Lett.* 8:1-7). Gamma-exotoxins are various proteolytic enzymes, chitinases and proteases. The toxic effects of gamma exotoxins are only expressed in combination with beta-exotoxin or delta-endotoxin. Forsberg *et al.*
15 (1976) "Bacillus thuringiensis: Its effects in Environmental Quality," National Research Council of Canada. Stonard *et al.* (1994) *ACS Symposium Series* 551: 25 report a water-soluble secondary metabolite active against corn rootworm in the supernatant of a *Bacillus cereus* strain.

There are no documented strains of *Bacillus* with both fungicidal and corn
20 rootworm activity. There are no known metabolites produced by *Bacillus subtilis* that are of less than 10,000 molecular weight and are extractable in a non-polar solvent.

Disclosure of the Invention

25 A novel antibiotic-producing and metabolite-producing strain of *Bacillus subtilis*, previously identified as *Bacillus megaterium*, is provided that exhibits broad fungicidal and bactericidal activity and also exhibits corn rootworm activity. Also provided is a novel metabolite from the novel *B. subtilis* with activity on corn rootworm. Also provided is a method of treating or protecting
30 plants from fungal and bacterial infections comprising the step of applying an

effective amount of the antibiotic-producing *Bacillus subtilis*. The antibiotic-producing *Bacillus subtilis* can be provided as a suspension in a whole broth culture or as an antibiotic-containing supernatant obtained from a whole broth culture of the antibiotic-producing strain of *Bacillus*. Also provided is a method of treating or protecting plant roots from corn rootworm infestations comprising the step of applying an effective amount of the novel metabolite-producing *Bacillus subtilis*. The novel metabolite-producing *Bacillus subtilis* can be provided as a suspension in a whole broth culture or as a metabolite-containing supernatant or a purified metabolite obtained from a whole broth culture of the microorganism. Also provided are novel compounds, agrastatins, produced by the novel strain AQ713 and a novel combination of compounds comprising inturin A, a plipastatin, a surfactin and an agrastatin.

Brief Description of the Drawings

Figure 1 shows the structure of the plipastatin antibiotics.
Figure 2 shows the HPLC chromatogram of AQ713 metabolites.

Modes of Carrying Out the Invention

The present invention provides a novel strain, AQ713, of *Bacillus subtilis*, previously identified as a *Bacillus megaterium*, or mutants thereof with the broad antifungal and antibacterial activity and the novel combination of antifungal and anti-corn rootworm activity. This novel strain is designated AQ713 and was deposited with the NRRL on March 7, 1997 under the provisions of the Budapest Treaty on the International Recognition of the Deposit of Microorganisms for the Purpose of Patent Procedure under Accession No. B21661. The invention also includes methods of preventing and treating fungal and bacterial diseases in plants using such bacterial strains or antibiotic-containing supernatants or pure antibiotics obtained from such bacterial strains. The invention also includes methods of treating plant roots or soil to control corn rootworm larvae with a bacterial suspension of AQ713 or a metabolite-containing supernatant of a culture

of AQ713 or purified metabolites from strain AQ713. The invention also includes a solvent-extractable metabolite with activity on corn rootworm with a molecular weight of less than 10,000 daltons. The invention further includes novel compounds, agrastatins, produced by the novel microorganism. Also included is a novel combination comprising an A-type iturin, a plipastatin, a surfactin and an agrastatin.

Definitions

As used herein, "biological control" is defined as control of a pathogen or insect by the use of a second organism. Known mechanisms of biological control include enteric bacteria that control root rot by out-competing fungi for space on the surface of the root. Bacterial toxins, such as antibiotics, have been used to control pathogens. The toxin can be isolated and applied directly to the plant or the bacterial species may administered so it produces the toxin *in situ*.

The term "fungus" or "fungi" includes a wide variety of nucleated spore-bearing organisms that are devoid of chlorophyll. Examples of fungi include yeasts, molds, mildews, rusts, and mushrooms.

The term "bacteria" includes any prokaryotic organism that does not have a distinct nucleus.

"Fungicidal" means the ability of a substance to increase mortality or inhibit the growth rate of fungi.

"Antibiotic" includes any substance that is able to kill or inhibit a microorganism. Antibiotics may be produced by a microorganism or by a synthetic process or semisynthetic process. The term, therefore, includes a substance that inhibits or kills fungi for example, zwittermicin-A or kanosamine.

"Antifungal" includes any substance that is able to kill or inhibit the growth of fungi.

The term "culturing" refers to the propagation of organisms on or in media of various kinds. "Whole broth culture" refers to a liquid culture containing both cells and media. "Supernatant" refers to the liquid broth remaining when cells

grown in broth are removed by centrifugation, filtration, sedimentation, or other means well known in the art.

5 An "effective amount" is an amount sufficient to effect beneficial or desired results. An effective amount can be administered in one or more administrations. In terms of treatment and protection, an "effective amount" is that amount sufficient to ameliorate, stabilize, reverse, slow or delay progression of the fungal or bacterial disease states.


10 As used herein, the term "insects" includes all organisms in the class "Insecta." "Pre-adult" insects refers to any form of an organism prior to the adult stage, including, for example, eggs, larvae, and nymphs. "Insecticidal" refers to the ability of a substance to increase mortality or inhibit growth rate of insects. "Nematicidal" refers to the ability of a substance to increase mortality or inhibit the growth rate of nematodes. "Pesticidal" refers to the ability of a substance to increase mortality or inhibit the growth rate of insects, nematodes and mites.

15 "Positive control" means a compound known to have pesticidal activity. "Positive controls" include, but are not limited to commercially available chemical pesticides. The term "negative control" means a compound known not to have pesticidal activity. Examples of negative controls are water or ethyl acetate.

20 The term "solvent" includes any liquid that holds another substance in solution. "Solvent extractable" refers to any compound that dissolves in a solvent and which then may be isolated from the solvent. Examples of solvents include, but are not limited to, organic solvents like ethyl acetate.

25 The term "metabolite" refers to any compound, substance or byproduct of a fermentation of a microorganism that has pesticidal activity. Antibiotic as defined above is a metabolite specifically active against a microorganism.

The term "agrastatins" refers to a group of novel compounds having the following structures:



where R₁ is a branched or straight aliphatic side chain, C8-C20; X is either
 5 Ala or Val; R₂ is an acetate or an ester derivative; and Glx is Gln or Glu. These
 compounds have antibacterial and antifungal activity as well as anti-corn
 rootworm activity.

We describe a novel metabolite and antibiotic-producing strain of *Bacillus*
subtilis, previously identified as *Bacillus megaterium*, that has broad antifungal
 10 and antibacterial activity and that also kills or stunts corn rootworm larvae. In
 another aspect, the present invention provides a method of treating or protecting
 plants from fungal and bacterial infections comprising applying an effective
 amount of a supernatant obtained from a whole broth culture of *Bacillus subtilis*
 AQ713 within the present invention. The supernatant may be obtained well
 15 known in the art including centrifugation, filtration, sedimentation and the like.

In another aspect, the invention encompasses a method of treating or
 protecting plants from fungal and bacterial infections comprising applying an
 effective amount of the whole broth of the novel strain *Bacillus subtilis*.

In further aspect, the invention encompasses a method of treating or
 20 protecting plants from fungal and bacterial diseases comprising applying an
 effective amount of the antibiotic produced by the novel strain of *Bacillus subtilis*.

In another aspect, the present invention provides a method of treating or
 protecting plant roots from corn rootworm infestations comprising applying an
 effective amount of a supernatant obtained from a whole broth culture of *Bacillus*
 25 *subtilis* AQ713 within the present invention. The supernatant may be obtained
 well known in the art including centrifugation, filtration, sedimentation and the
 like.

In another aspect, the invention encompasses a method of treating or
 protecting plants from corn rootworm infestations comprising applying an
 30 effective amount of the whole broth of the novel strain *Bacillus subtilis*.

In further aspect, the invention encompasses a method of treating or protecting plant roots from corn rootworm infestations comprising applying an effective amount of the metabolite produced by the novel strain of *Bacillus subtilis*.

5 In order to achieve good dispersion and adhesion of compositions within the present invention, it may be advantageous to formulate the whole broth culture, supernatant and/or metabolite/antibiotic with components that aid dispersion and adhesion. Suitable formulations will be known to those skilled in the art.

10 Compositions within the present invention can be formulated as wettable powders, granules and the like, or can be microencapsulated in a suitable medium and the like. Examples of other formulations include, but are not limited to soluble powders, wettable granules, dry flowables, aqueous flowables, wettable dispersible granules, emulsifiable concentrates and aqueous suspensions. Other
15 suitable formulations will be known to those skilled in the art.

 In yet a further aspect of the present invention, a novel group of compounds designated "agrastatins" are provided. These compounds exhibit antibacterial and antifungal activity in addition to anti-corn rootworm activity.

 In still a further aspect of the present invention, a novel combination
20 comprising an A-type iturin, a plipastatin, a surfactin and an agrastatin is provided.

 In another aspect of the present invention, methods of treating or protecting plants from fungal and bacterial diseases comprising applying an effective amount of a novel combination of compounds comprising an A-type
25 iturin, a plipastatin, a surfactin and an agrastatin are provided.

 All patents and publications cited herein are hereby incorporated by reference in their entirety. The following examples are provided to illustrate the invention. These examples are not to be construed as limiting.

EXAMPLES

Example 1

Characterization of Strain AQ713

5 The isolate was identified based on utilization of the Biolog microplate panel (Biolog, Inc., Hayward, CA) as described in Bochner (1989) *Nature* 339: 157-158. The Biolog microplate is comprised of prefilled and dried panel wells with 95 different carbon substrates plates available for gram positive and gram negative bacteria. The isolate was grown in liquid medium at 28°C and after 24 hrs a washed cell suspension (0.85% saline) was inoculated into each panel well
10 of a GP Microplate (Biolog, Inc.) After 24 hrs at 28°C, carbon utilization reactions were assessed. Substrate utilization profiles were then compared to the Biolog Gram-Positive Data Base (release 3.50) and isolated to closest similar species. Biolog results gave a similarity index of 0.883 to *Bacillus megaterium*.

15 A more extensive characterization of AQ713 was conducted by the American Type Culture Collection, Rockville, Md.

Isolate submitted as: Unknown; Strain AQ 713

Isolate identified as: Using the available physiological and biochemical data, this strain most closely resembles *Bacillus subtilis*.

Cellular morphology: The motile cells are found in singly, with one endospore formed in the central or subterminal region. The cells are uniformly stained Gram positive.

Colonial morphology: The colonies are opaque and irregular with convex elevation, a rough, dull surface and an erose margin.

Characterization Data of Strain AQ 713:

Rods	+	Colony opaque	+
Rods straight	+	Colony entire	-
Rods curved	-	Colony erose	+
Cells single	+	Colony lobate	-
Cells chained	-	Colony circular	-
Ends tapered	-	Colony irregular	+
Ends rounded	+	Colony rhizoid	-
Ends squared	-	Colony low convex	+
Endospore formed	+	Colony high convex	-
Sporangium swollen	-	Colony flat	-
One spore/cell	+	Colony raised	-
Spore round	-	Colony glistening	-
Spore cylindrical	+	Colony dull	+
Spore oval	+	Colony dry	-
Spore central	+	Colony smooth	-
Spore terminal	-	Colony rough	+
Spore subterminal	+	Soluble brown pigment	-
Gram stained	+	Soluble black pigment	-
Gram positive	+	Soluble yellow pigment	-
Gram negative	-	Insoluble brown pigment	-
Gram variable	-	Insoluble black pigment	-
Vacuoles present	-	Insoluble yellow pigment	-
Colony translucent	-	Insoluble orange pigment	-
Colony transparent	-	Insoluble red pigment	-

Cells motile	+
Growth at 15°C	+
Growth at 20°C	+
Growth at 26°C	+
Growth at 30°C	+
Growth at 37°C	+
Growth at 45°C	+
Growth at 50°C	weak
Growth at 55°C	-
Growth at 60°C	-
Growth at 65°C	-
Catalase	+
Oxidase	+
Casein hydrolysis	+
Gelatin liquification	+
Hippurate hydrolysis	-
Lecithinase degradation	-
Starch hydrolysis	+
Tween 80 hydrolysis	+
Tyrosine decomposition	-
Growth in 2% NaCl	+
Growth in 5% NaCl	+
Growth in 7% NaCl	+
Growth in 10% NaCl	+
Growth in 0.2% Na azide	V
Growth at pH 4.5	+
Growth at pH 6.0	+
Acid from arabinose	-
Gas from arabinose	-
Acid from cellobiose	weak
Acid delayed > 14 days	weak
Gas from cellobiose	-
Acid from fructose	+
Acid delayed > 14 days	-
Gas from fructose	+
Acid from glucose	+
Acid delayed > 14 days	-
Gas from glucose	-

Acid from lactose	-
Gas from lactose	-
Acid from mannitol	-
Gas from mannitol	-
Acid from mannose	-
Gas from mannose	-
Acid from sucrose	weak
Acid delayed >14 days	weak
Gas from sucrose	-
Acid from trehalose	-
Gas from trehalose	-
Acid from xylose	-
Gas from xylose	-
Aerobe	-
Faculative	-
Microaerophile	+
Anaerobe	-
Gas from sealed nitrate	-
Gas from sealed glucose	-
Indole	-
Nitrate to nitrite	+
Nitrate to gas	-
Methylene blue reduction	+
Methylene blue reoxidation	-
Litmus milk acid	-
Litmus milk coagulated	-
Litmus milk alkaline	+
Litmus milk reduced	+
Litmus Milk peptonized	+
VP (5198) positive	+
VP (5331) positive	+
pH VP 5198 6.0 or less	-
pH VP 5198 6.5 - 7.5	+
pH VP 5198 8.0 or more	-
Citrate utilization	+
Propionate utilization	-
Propionate utilization	-

Comments: Using the available physiological and biochemical data, this strain most closely resembles *Bacillus subtilis*.

Key Characterization Results

Characterization Tests	Strain AQ 713	<i>Bacillus subtilis</i>
Swollen sporangium	-	-
Anaerobic growth	microaerophilic	microaerophilic
VP reaction	+	+
pH of VP	7.0	5.0 - 8.0
Maximum temperature growth	55°C	45 - 55°C
7% NaCl growth	+	+
Acid from glucose	+	+
Acid from arabinose	-	+
Acid from xylose	-	+
Acid from mannitol	-	+
Casein decomposition	+	+
Tyrosine decomposition	-	-
Citrate utilization	+	+
Propionate utilization	-	-

Reference:

Gordon, R.E., W.C. Haynes and C.H.N. Pang. 1973. The Genus Bacillus. Handbook No. 427. U.S. Department of Agriculture, Washington, D.C.

Example 2

Activity of AQ713 Against Corn Rootworm

Bacillus samples were grown in a *Bacillus* culture media. Medium 2 contained 5% peptone, 5% dextrose, 3% yeast extract, 3% malt extract, 1.5%
5 proflo cotton seed extract (59% protein, 4.26% fat, 6.73% ash, 3.19% fiber and trace amounts of gossypol; the balance is water), 10% soy flour, and 0.5 % $\text{MgSO}_4 \times 7\text{H}_2\text{O}$. Medium 3 contained the same ingredients, except with 20% peptone and 3.4% KH_2PO_4 and 4.3% K_2HPO_4 . One day old streaked cultures were used to inoculate 250 mL baffled shake flasks. Flasks were shaken at 200 rpm at
10 29°C for 5 days. To assay insecticidal activity, 35 mL of culture broth were centrifuged at 5,200 rpm for 20 minutes and the supernatant used in microassay described below.

Assays were performed in 96-well microplates. Each well contained a solid agar substrate, a test organism and either a positive control, a negative
15 control or supernatant obtained as described in Example 1 from the novel *Bacillus* strain.

To assay insecticidal activity, an agar substrate was prepared for the wells of the microplate according to Marrone *et al.* (1985), *J. Econ. Entomol.* 78: 290-293. To assay nematicidal activity, plain agar (1.5%) was used in the wells
20 instead.

A 1 ppm solution of Avid® (avermectin) was used as a positive control. Deionized water was used as a negative control. Two replicates of test sample or control were used for each assay. 40 uL of supernatant sample or whole broth grown in medium 1, 2 or 3 were dispensed into each sample well. Plates were then
25 placed in a fume hood to dry for approximately 2-3 hours until the agar solution was dried.

Test organisms were either pre-adult corn rootworms (*Diabrotica undecimpunctata*), pre-adult German cockroaches (*Blatella germanica*), pre-adult beet armyworms (*Spodoptera exigua*), pre-adult flies (*Drosophila melanogaster*),
30 or the N2 strain of the nematode *Caenorhabditis elegans*. Test organisms were

diluted in 0.1% agar to a concentration of approximately 5 organisms per 25 uL of agar dispensed into each well. The microplate was sealed with an airtight substance such as Mylar®, and each well ventilated with a pin press. The plates were incubated at 27°C for up to 7 days.

5 After incubation, wells were scored by noting neonate mortality or the degree of larval development. Sample wells containing all dead or stunted larvae were given a score of 1, wells containing some dead and other severely stunted larvae were given a score of 2, live but stunted larvae were scored as 3 and sample wells containing no dead larvae were given a score of 4. Scores were averaged
10 among replicates within each sample. Results are summarized in Tables 2 and 3.

Table 2: Score Rating of AQ713 Against Insect Pests Whole broth

	<i>C. elegans</i>	Corn rootworm	Beet armyworm	Fruit Fly	Positive Control	Negative Control
Medium 2	NT	1.0	4.0	4.0	1.0	4.0
15 Medium 3	NT	2.0	4.0	4.0	1.0	4.0

NT = not tested

Table 3: Score Rating of AQ713 Against Insect Pests Supernatant

	<i>C. elegans</i>	Corn rootworm	Beet armyworm	Fruit Fly	German Cockroach	Positive Control	Negative Control
20 Medium 2	4.0	3.0	4.0	4.0	4.0	1.0	4.0
Medium 2	4.0	4.0	4.0	4.0	4.0	1.0	4.0

These tests show that AQ713 was active in both media as a whole broth culture, with the best activity in medium 2. The supernatant was only active when AQ713 was grown in medium 2.

Example 3**Chemical Properties of the AQ713 Metabolite****Active Against Corn Rootworm**

50 mL of AQ713 was grown in media 2. To each culture was added 50
5 mL ethyl acetate and the mixture was shaken in a separatory funnel for 2 minutes.
The aqueous layer was removed and the organic layer was collected in a bottle
containing magnesium sulfate. The organic filtrate was then filtered into a round
bottom flask and the solvent removed on the rotovap.

For the bioassay, the dried organic extract was redissolved in 2.5 mL
10 acetone. A 40 uL aliquot was removed and diluted to 800 uL with 70%
acetone/water. This is a 10X concentration of the organic extract. Serial dilutions
were carried out to obtain samples on neonate corn rootworm with percent
mortality recorded of neonate larvae (1 per well in a microtiter plate as prepared
above) after 7 days. The results are recorded in Table 4.

15 **Table 4: Activity of Ethyl Acetate Extracts of AQ713 Against Corn Rootworm**

<u>Sample</u>		<u>Percent Mortality</u>
AQ713:	Organic extract 10X	89
	Organic extract 5X	93
	Organic extract 1X	65
	Whole broth	100
	70% acetone/water	27
	Water	59

25 The results show that AQ713 produces a solvent-extractable metabolite that kills
corn rootworms.

To determine the molecular weight range of the active metabolite, a 50-mL
culture of AQ713 was grown in media 2. One mL was placed into a microfuge
tube and spun at 12,000 rpm for 15 minutes. The supernatant was removed. 500
30 microliters of supernatant was placed on top of a 10,000 dalton molecular weight

centricon filter. These were centrifuged according to the manufacturer's instructions (12,000 rpm for 35 minutes). The filtrate was collected and the retentate recovered by centrifugation and washing of the filter. Samples of the supernatant, filtrate and retentate were tested against neonate corn rootworm larvae (96 well-plate with insect diet, Marrone *et al.*, *supra* as above; 40 uL of sample per well and 8 wells for each sample, 1 larva/well). The results of the test are shown in Table 5.

Table 5: Molecular Weight Cutoff of AQ713

10

Percent Mortality
Against Corn Rootworm

15

AQ713:	supernatant	43
	filtrate	63
	retentate	17

The results show that the supernatant and filtrate were active, thus the molecular weight of the metabolite is less than 10,000 daltons.

20

Example 4

Chemical properties of the AQ713 metabolite active against plant pathogens

25

50 mL of AQ713 was grown in media 2. To each culture was added 50 mL ethyl acetate and the mixture was shaken in a separatory funnel for 2 minutes. The aqueous layer was removed and the organic layer was collected in a bottle containing magnesium sulfate. The organic filtrate was then filtered into a round bottom flask and the solvent removed on the rotovap.

For the bioassay, the dried organic extract was redissolved in 2.5 mL acetone. A 40 uL aliquot was removed and diluted to 800 uL with 70% acetone/water. This is a 10X concentration of the organic extract. A 96-well plate

assay (described below) plant pathogen assay with *Pythium ultimum* and *Botrytis cinerea* was conducted to determine activity of the organic extract. The whole broth gave 100% control (score of 1), but the 10X organic extract gave no control of the two plant pathogens (score of 4). This indicates that the active antibiotics, unlike the corn rootworm active metabolites produced by AQ713 are not extractable in an organic solvent such as ethyl acetate.


Further testing provided for the isolation of a novel compound, agrastatin A. A butanol extract was made of the fermentation broth by first extracting the broth two times with an equal volume of ethyl acetate and separating the layers. The aqueous fraction was then extracted two times with an equal volume of butanol. The butanol extracts were combined and solvent was removed with a rotary evaporator. A powder was obtained by freeze drying the resulting extract.

The powder was dissolved in 80% acetonitrile/water and sonicated. The solution was applied to a C-18 solid phase extraction (SPE) cartridge that had been activated with methanol and equilibrated with 80% acetonitrile/water. The SPE cartridge was eluted with 80% ACN/water and this eluent was collected and the solvents removed. The eluent was further purified by HPLC. A C-18 HPLC column (1 cm X 25 cm) was used (UV detection at 210 nm) with an acetonitrile + 0.05% TFA/water + 0.05% TFA solvent gradient as follows: 0-20 minutes, 33% ACN; 20-30 minutes, 40% ACN; 30-45 minutes, 45-55% ACN; and 45-63 minutes, 55% ACN.

An HPLC chromatogram of AQ713 shows the presence of the iturins, iturin-like compounds (plipastatins and agrastatins) and surfactins, see Figure 1. Iturins A2, A3, A4, A7 and A6 were identified by a combination of NMR data and LC mass spectrometry data and comparison to literature values. Surfactins were identified by comparison to purchased surfactin standards by HPLC and by LC mass spectrometry.


The iturin-like compounds were determined to be a mixture of plipastatins and the novel agrastatins by a combination of amino acid analysis and LC mass spectrometry. Extensive NMR data was also collected for one of the novel

compounds (HPLC peak 20), designated agrastatin A. Agrastatin A was found to contain the following amino acids: Thr; 3 Glu; Pro; Ala; Val; 2 Tyr; and Orn. This make up differs from plipastatin A by the presence of Val and the loss of Ile. The molecular weight of agrastatin A was determined to be 1448 which
 5 corresponds to the following structure:



The straight chain nature of the fatty acid portion was confirmed by ¹H
 10 NMR. The position of the amino acids in the cyclic peptide was determined by detailed analysis of the TOCSY and ROESY datasets.

Mass spectrometry and amino acid analysis of agrastatin B (HPLC peak 26) suggest that its structure is similar to plipastatin B2 with the substitution of the Ala residue with Val. The structure is shown below:



Example 5

20 Activity of AQ713 Against Plant Pathogens in *in-vitro* Culture (96-well plate)

To determine if AQ713 is effective against the fungi, *Phytophthora infestans*, *Pythium ultimum*, *Botrytis cinerea*, *Rhizoctonia solani*, *Alternaria solani*, the following experiments were performed. 96-well plates (flat-bottomed, 400 microliters per well, Nunc brand) were filled with an agar medium (potato dextrose agar) (PDA, Difco). *Phytophthora infestans* cultures were grown for
 25 three days in liquid YPG-1 medium (0.4 g yeast, 0.1% KH₂PO₄, 0.5% MgSO₄ X 7 H₂O, 1.5 % glucose). For the other fungi, spores were scraped from the surface of petri plates and 0.1-0.2 mL aliquots of deionized water and spore suspension (concentration approximately 2 X 10⁶ spores/mL) of pathogen were spread onto
 30 the agar.

AQ713 was grown for 72 hours in the medium 2 or 3 as described in Example 2. To obtain supernatants, the whole broth culture was centrifuged at 5,200 rpm for 20 minutes. The fungal plant pathogens were pipetted onto the 96-well plates (8 wells/pathogen). The presence or absence of fungal growth was recorded for each of 8 wells. Approximately 40 uL of AQ713 supernatant or 20 uL of whole broth was added to each well. A score of "1" means complete inhibition of fungal growth. A score of "4" means no inhibition of fungal growth. Results are shown in Table 6.

Table 6: *In-Vitro* Inhibition of Fungal Growth (96-well plate)

	<u>AQ713 Supernatant</u>	<u>Media 2</u>	<u>Media 3</u>
		<u>Score</u>	<u>Score</u>
15	<i>Phytophthora infestans</i>	1	1
	<i>Pythium ultimum</i>	1	1
	<i>Botrytis cinerea</i>	1	1
	<i>Rhizoctonia solani</i>	4	1
	<i>Alternaria solani</i>	1	1
25	<u>AQ713 Whole broth</u>		
	<i>Colletotrichum cocodes</i>	1	NT
	<i>Alternaria brassicicola</i>	1	NT
	<i>Botrytis cinerea</i>	1	NT
	<i>Cladosporium cucumerinum</i>	1	NT
	<i>Monilinia fructicola</i>	1	NT
	<i>Venturia pyrina</i>	1	NT
	<i>Rhizoctonia solani</i>	1	NT
	<i>Alternaria solani</i>	1	NT
30	NT Not tested		

The results show that AQ713 has broad fungicidal spectrum *in-vitro* and that both the whole broth and supernatant are highly active. The supernatant was active on *Rhizoctonia solani* in medium 3 but not medium 2.

Example 6

Activity of AQ713 Against Plant Pathogens in *in-vitro* Culture (zone assay)

To determine the activity of AQ713 in an agar diffusion (zone) assay, plant pathogen spores were spread over the surface of potato dextrose agar in 10 cm petri dishes. 7.0 mm wells were removed from the agar and a 100 uL sample of the supernatant of AQ713 grown in medium 2 was placed in the well. Supernatant was prepared by centrifuging at 4200 rpm for 40 minutes. The supernatant was then spun again at 4200 rpm for another 40 minutes. Typical results consisted of a zone of no growth and/or reduced growth of the pathogen around the well. The zone size in millimeters was measured and recorded. The results are shown in Table 7.

Table 7: *In-Vitro* Inhibition of Fungal Plant Pathogen Growth (Zone Test)

	<i>Alternaria brassicicola</i>	<i>Botrytis cinerea</i>	<i>Monilinia fructicola</i>
AQ713 supernatant Zone size (mm)	16	23	14
AQ713 Whole broth	22	15	18

Example 7

Activity of AQ713 Against Bacterial Plant Pathogens

A standard agar diffusion assay was set up as in example 6. A lawn of each bacterial pathogen was spread over the surface of a petri plate. 100 uL of AQ713 whole broth grown in medium 2 was placed in each well. The size of the zone was measured in millimeters.

Table 8: *In-Vitro* Inhibition of Bacterial Plant Pathogens (Zone Test)

AQ713 Whole broth:	Inhibition Zone (mm)
<i>Acidovorax avenae</i> subsp. <i>citrulli</i>	18
<i>Pseudomonas syringae</i> pv. <i>tomato</i>	11
5 <i>Xanthomonas campestris</i> pv. <i>campestris</i>	18
<i>Erwinia carotovora</i> subsp. <i>carotovora</i>	11
<i>Clavibacter michiganense</i> subsp. <i>michiganense</i>	22

AQ713 was active against all species of bacterial plant pathogens tested *in-vitro*.

10

Example 8**Activity of AQ713 Against Plant Pathogens in Plant Tests**

The activity of AQ713 was tested against gray mold, *Botrytis cinerea*, on beans and geranium leaves, *Alternaria solani* on tomato seedlings, and downy mildew of lettuce, *Bremia lactucae*.

15

For *A. solani*, tomato seedlings at the 2-3 leaf stage planted in 6-packs were sprayed to runoff with AQ713 whole broth (media 2). After spraying, the seedlings were allowed to dry (about 1.5 hours). The seedlings were then sprayed with 5.0×10^4 spores/mL. Seedlings were covered with a plastic dome and kept at 28°C in a Percival incubator. Water with no AQ713, with and without spores of the pathogen was used as a negative control and a positive pathogen control. Four days later the test was read. On the water *A. solani* control, there were uniform lesions over all the leaves and the cotyledons were detached and severely infected (rating of 5 = complete infection, no control). AQ713 treated plants had a few light lesions scattered on the true leaves. The cotyledons were attached but with some small lesions (rating of 1). The negative control was not infected.

20

25

A second test was set up using detached tomato seedlings (stems broken off at the ground level) placed in mason jars filled with water put under domes and stored as above. The plants were sprayed as above and the symptoms of *A.*

solani were recorded four days later. There were no symptoms on the negative control. On the positive control, there were uniform lesions over the seedlings. The AQ713 treatment was rated 1 (few or no lesions). Two days later, the plants in the positive control were destroyed, but the AQ713 treated seedlings were
5 virtually clean and looked the same as the negative controls (water sprayed plants).

For the test on *Botrytis cinerea*, the first true leaves of a bean plant were wounded by pressing the mouth of a 13 X 100 culture tube onto each leaf. Each leaf received two wounds/leaf. The leaves were sprayed with AQ713 whole broth
10 (media 2) or water alone or the pathogen alone. When dry, they were again sprayed with *B. cinerea* spores (0.8×10^6 spores/mL). The leaves were placed in flats covered with plastic domes and stored at 18-20°C in a Percival incubator. Five days later, the positive control (pathogen alone) was rotted in an area about 25 mm in diameter. The negative control (water alone) had no rotting. AQ713
15 showed no infections on 7 of 8 circles where the leaves were wounded. The one that was infected had light infection at two locations around the circle.

For the *Bremia* test, lettuce seeds were planted in a layer of sterilized potting mix containing peat, perlite and vermiculite in small clear plastic plant condominiums about 8 centimeters high and wide. After the lettuce germinated
20 (one week), the lettuce seedlings were sprayed with the AQ713 broth or supernatant sample. The plants were allowed to dry and then downy mildew spore suspension from infected lettuce seedlings was sprayed onto the seedlings. The plastic covers were placed over the plants and incubated at 18-20 20°C in a Percival incubator. One week later, the test was evaluated. AQ713 did not
25 prevent downy mildew from *Bremia* on lettuce seedlings.

Example 9

Efficacy of AQ713 Against Plant Diseases (Greenhouse Test)

Grape Downy Mildew

AQ713 was grown in a soy-based medium in a 400 liter fermenter for 48
5 hours. Grape plants (cultivar Chardonnay) were sprayed with a hand-held sprayer
to run-off with whole broth from the 400 liter fermentation run diluted with sterile
water to 0.5X and 0.25 X concentrations. When the foliage dried, the plants were
sprayed a second time. After drying, the plants were inoculated with the pathogen
causing grape downy mildew, *Plasmopara viticola*. Three plants were treated for
10 each dose. Each plant was evaluated by estimating the percent disease control
based on a scale from 0 to 100% control. 100% control is a plant with no visible
lesions. A chemical fungicide, metalaxyl, was used for comparison. The results
were as follows:

	AQ713 0.5X whole broth	97.7% control
15	AQ713 0.25X whole broth	100% control
	Metalaxyl 30 ppm	100% control
	Metalaxyl 10 ppm	98.3% control
	Metalaxyl 1 ppm	80% control

The results demonstrate that AQ713 effected control of grape downy mildew as
20 well as the chemical fungicide.

Example 10

Efficacy of AQ713 Against Squash Powdery Mildew

AQ713 was grown in a soy-based medium in a 400 liter fermenter for 48
25 hours. Squash plants (Crookneck and Acorn) were sprayed with a hand-held
sprayer to run-off with whole broth from the 400 liter fermentation run and a
sample diluted with sterile water to 0.5X concentration. After drying, the plants
were inoculated with the squash powdery mildew pathogen, *Sphaerotheca*
fuliginea. Two plants were treated for each dose. Spray dried powder of the
30 whole broth was also tested. The 400 liter fermentation broth was spray dried to

remove the water. 10% and 2.5% spray dried powder solutions were sprayed on the plants to run-off as above. The incidence of powdery mildew disease was rated on a score from 0 to 5. The 5 rating is 100% disease whereas the 0 rating is no disease. The results are shown below in Table 9.

5

Table 9

Test Suspension	Acorn Squash Plant 1	Acorn Squash Plant 2	Crookneck Squash Plant 1	Crookneck Squash Plant 2
AQ713 1X whole broth	0	0	0	0
AQ713 0.5X whole broth	0	0	0	0
AQ713 10% spray dried powder	0	0	0	0
AQ713 2.5% spray dried powder	0	0	0.5	1

AQ713 whole broth and spray dried powder provided nearly complete control of squash powdery mildew.

10

Example 11

Efficacy of AQ713 on Late Blight, Gray Mold, Grape Powdery Mildew, Cereal Powdery Mildew, Sheath Blight and Rice Blast in the Greenhouse

AQ713 was grown in a soy-based medium for 72 hours in a 250 mL shake flask. The disease, causative pathogen and host are listed in Table 10 below. This whole broth culture was tested on the plants as shown in Table 11 below.

15

Table 10

Disease	Plant Pathogen	Host
Late Blight	<i>Phytophthora infestans</i>	Tomato
Gray Mold	<i>Botrytis cinerea</i>	Pepper
Sheath Blight	<i>Rhizoctonia solani</i>	Rice
Rice Blast	<i>Pyricularia oryzae</i>	Rice
Powdery Mildew	<i>Uncinula necator</i>	Grape
Powdery Mildew	<i>Drysiphe graminis</i> f. sp. <i>graminis</i>	Wheat

5 Each broth was sprayed to run-off at 1X concentration on the test plants with a hand held sprayer, allowed to dry and then sprayed a second time. Three plants were treated for each disease and treatment. After drying, the plants were inoculated with the pathogens. Each plant was evaluated by estimating the percent disease control based on a scale from 0 to 100% control 100% control refers to a plant with no visible lesions. Chemical fungicides were used for comparison. Disease index is the severity of the disease on the untreated control.

Table 11

	<i>P. infestans</i>	<i>B. cinerea</i>	<i>E. graminis</i>	<i>U. necator</i>	<i>P. oryzae</i>	<i>R. solani</i>
AQ713	70	100	84	100	100	100
Metalaxy 30 ppm	100					
Metalaxyl 10 ppm	77					
Propico- nazole 10 ppm		87				
Propico- nazole 5 ppm		57				
Propico- nazole 0.5 ppm			100			
Propico- nazole 0.2 ppm			54			
Myclo- butanil 30 ppm				100		
Myclo- butanil 10 ppm				100		
Pencycuron 50 ppm					100	
Pencycuron 10 ppm					100	
Benomyl 100 ppm						100
Benomyl 40 ppm						77
Disease Index (%)	80	95	70	50	60	80

AQ713 showed activity that was equivalent to chemical fungicides on all the pathogens tested.

Example 12

5 **Efficacy of AQ713 Against *Brassica* Downy Mildew**

10 *Bacillus* strain AQ713 was grown in a ten liter fermenter in a soy-based medium for 48 hours. The whole broth culture at 1X strength was sprayed onto three week-old cauliflower and brussel sprouts plants at the full cotyledon stage with an artist's air brush powered by compressed air. Three replicates of 15-25 seedlings/pot were sprayed per treatment. Quadris™, an azoxystrobin fungicide from Zeneca, was also sprayed on plants (three per treatment) at rates of 250 ppm and 125 ppm. A spore suspension of downy mildew, *Peronospora parasitica*, at 1-5 X 10⁴ spores/mL was sprayed onto the *Brassica* plants after the AQ713 and Quadris sprays dried. The plants were held at 15-17°C for 24 hours for infection, 15 then the seedlings were incubated at 20-24°C for six days. The pots were returned to 15-17°C overnight to allow sporulation of the pathogen until the test was rated. Each plant was evaluated by estimating the percent disease control based on a scale from 0 to 100% control. 100% control is a plant with no sporulating lesions. The results averaged across replicate pots are shown below in Table 12.

20

Table 12

	Reading taken December 23	Reading taken December 30	Reading taken January 6
AQ713 whole broth	100	90	75
Quadris 250 ppm	100	NT	NT
Quadris 125 ppm	NT	100	100
Water Control	0	0	0

NT = Not Tested

AQ713 controlled downy mildew effectively for three weeks in duration.

Example 13

Synergism of AQ713 and a Commercial Fungicide

AQ713 was grown in a ten liter fermenter in a soy-based medium for 72 hours. The bacterial culture was diluted with sterile water to 0.5X and 0.25X concentrations. The culture at 1X, 0.5X and 0.25X concentrations was sprayed onto three week-old pepper plants with an artist's air brush powered by compressed air. Three plants were sprayed per treatment. Quadris™, an azoxystrobin fungicide from Zeneca, was also sprayed on plants (three per treatment) at concentrations of 500 ppm, 250 ppm and 125 ppm. In addition, combinations of Quadris plus the whole broth culture of AQ713 in a 1:1 ratio were sprayed onto pepper plants (three per treatment). The treatments with and without Quadris are outlined in Table 13 below. A spore suspension of *Botrytis cinerea*, gray mold, at 1×10^6 spores/mL was sprayed onto the pepper plants after the AQ713 and Quadris sprays dried. The plants were held at 20-22°C for 3 days until the test was rated. The incidence of gray mold disease was rated on a score from 0 to 5. The 5 rating indicates 100% disease whereas the 0 rating indicates no disease. The results are shown in Table 13 below.

Table 13

Treatment	Rating Replicate 1	Rating Replicate 2	Rating Replicate 3	Rating Average
AQ713 1X	0.5	0.5	1.5	0.8
AQ713 0.5X	2.0	2.5	2.0	2.2
AQ713 0.25X	3.0	3.0	2.0	2.7
Quadris 500 ppm	4.0	3.5	4.0	3.8
Quadris 250 ppm	2.5	3.5	3.0	3.0
AQ713 1X + Quadris 500 ppm	0.5	1.0	1.0	0.8
AQ713 1X + Quadris 250 ppm	1.0	1.0	0.5	0.8
AQ713 0.5X + Quadris 250 ppm	0.5	1.0	1.0	0.8
AQ713 0.25X + Quadris 250 ppm	0.5	1.0	2.5	1.3
Water control	4.0	5.0	5.0	4.7
Water control 2	5.0	5.0	5.0	5.0

The results clearly show that combinations of Quadris and AQ713 control gray mold disease significantly better than either Quadris or AQ713 alone.

CLAIMS

What is claimed is:

1. An isolated, pure culture of *Bacillus subtilis* strain AQ713, NRRL
5 Accession No. B21661 and its mutants.
2. A metabolite produced by the *Bacillus subtilis* strain of claim 1
that exhibits activity against corn rootworm, is solvent extractable and has a
molecular weight of less than 10,000 daltons.
3. A supernatant obtained from a culture of the *Bacillus subtilis* strain
10 AQ713 of claim 1 that exhibits antifungal and antibacterial activity and activity
against corn rootworm.
4. A composition comprising the whole broth culture of the *Bacillus*
subtilis strain AQ713 of claim 1 and a chemical fungicide.
5. A composition comprising the whole broth culture of the *Bacillus*
15 *subtilis* strain AQ713 of claim 1 and a biological or chemical pesticide.
6. The composition of claim 5 further comprising a chemical
fungicide.
7. A composition comprising the metabolite of claim 2 and a
chemical fungicide.
8. A composition comprising the metabolite of claim 2 and a
20 biological or chemical pesticide.
9. The composition of claim 8 further comprising chemical fungicide.
10. A composition comprising the supernatant of claim 3 and a
chemical fungicide.
11. A composition comprising the supernatant of claim 3 and a
25 biological or chemical pesticide.
12. The composition of claim 11 further comprising a chemical
pesticide.

13. A method for protecting or treating plants and fruit from fungal and bacterial infections and corn rootworm infestations comprising applying an effective amount of the *Bacillus subtilis* strain of claim 1.

14. A method for protecting or treating plants and fruit from fungal and bacterial infections and corn rootworm infestations comprising applying an effective amount of the metabolite of claim 2.

15. A method for protecting or treating plants and fruit from fungal and bacterial infections and corn rootworm infestations comprising applying an effective amount of the supernatant of claim 3.

16. A method for protecting or treating plants and fruit from fungal and bacterial infections and corn rootworm infestations comprising applying an effective amount of the composition of claim 4, 5, 6, 7, 8, 9, 10, 11, or 12.

17. The method of claim 13, 14 or 15 wherein the infections are caused by at least one microorganism selected from the group consisting of *Phytophthora infestans*, *Rhizoctonia solani*, *Pythium ultimum*, *Botrytis cinerea*, *Alternaria solani*, *Colletotrichum cocodes*, *Alternaria brassicicola*, *Cladosporium cucumerinum*, *Monilinia fructicola*, *Venturia pyrina*, *Acidovorax avenae*, *Pseudomonas syringae*, *Xanthomonas campestris*, *Erwinia carotovora*, *Clavibacter michiganense*, *Plasmopara viticola*, *Sphaerotheca fuliginea*, *Uncinula necator*, and *Peronospora parasitica*.

18. The method of claim 16 wherein the infections are caused by at least one microorganism selected from the group consisting of *Phytophthora infestans*, *Rhizoctonia solani*, *Pythium ultimum*, *Botrytis cinerea*, *Alternaria solani*, *Colletotrichum cocodes*, *Alternaria brassicicola*, *Cladosporium cucumerinum*, *Monilinia fructicola*, *Venturia pyrina*, *Acidovorax avenae*, *Pseudomonas syringae*, *Xanthomonas campestris*, *Erwinia carotovora*, *Clavibacter michiganense*, *Plasmopara viticola*, *Sphaerotheca fuliginea*, *Uncinula necator*, and *Peronospora parasitica*.

19. The method of claim 13 wherein the *Bacillus subtilis* strain AQ713 is applied as a whole broth culture.

20. The method of claim 13, wherein the *Bacillus subtilis* strain AQ713 is applied as a supernatant.


21. The method of claim 19 wherein the *Bacillus subtilis* strain AQ713 is applied as wettable powders, granules, flowables or microencapsulations.

5 22. The method of claim 20, wherein the *Bacillus subtilis* strain AQ713 is applied as wettable powders, granules, flowables or microencapsulations.

23. The method of claims 13, 14 or 15 wherein the roots of plants or the soil around the roots are treated.


10 24. The method of claim 16 wherein the roots of plants or the soil around the roots are treated.

25. A compound having the formula




wherein R_1 is a branched or straight aliphatic side chain of C8-C20; X is either Ala or Val, R_2 is an acetate or an ester derivative and Glx is Gln or Glu.

26. A compound having the formula



27. A compound having the formula




28. A composition comprising an A-type iturin, a plipastatin, and a surfactant.

29. The composition of claim 28 further comprising an agrastatin.

30. A method for protecting or treating plants and fruit from fungal and bacterial infections and corn root worm infestations comprising applying an effective amount of the composition of claim 23, 24, 25, 26 or 27.

31. The method of claim 30 wherein the infections are caused by at least one microorganism selected from the group consisting of *Phytophthora infestans*, *Rhizoctonia solani*, *Pythium ultimum*, *Botrytis cinerea*, *Alternaria solani*, *Colletotrichum cocodes*, *Alternaria brassicicola*, *Cladosporium cucumerinum*, *Monilinia fructicola*, *Venturia pyrina*, *Acidovorax avenae*, *Pseudomonas syringae*, *Xanthomonas campestris*, *Erwinia carotovora*, *Clavibacter michiganense*, *Plasmopara viticola*, *Sphaerotheca fuliginea*, *Uncinula necator*, and *Peronospora parasitica*.

30. A method for protecting or treating plants corn rootworm infestations comprising applying an effective amount of the composition of claim 23, 24, 25, or 26 to roots of plants or to soil around the roots.



Plipastatin A1, X = Ala, R = CH₃(CH₂)₁₂-

Plipastatin A2, X = Ala, R = CH₃CH₂CHCH₃(CH₂)₁₀-

Plipastatin B1, X = Val, R = CH₃(CH₂)₁₂-

Plipastatin B2, X = Val, R = CH₃CH₂CHCH₃(CH₂)₁₀-

FIG. 1




FIG. 2

INTERNATIONAL SEARCH REPORT

International Application No.

PCT/US 98/09471

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C07K7/06 A01N63/02 C12N1/20 //(C12N1/20,C12R1:125)

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07K A01N C12N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P, X	<p>KIMURA, KEN-ICHI ET AL: "SNA-60-367, new peptide enzyme inhibitors against aromatase"</p> <p>J. ANTIBIOT. (1997), 50(6), 529-531 CODEN: JANTAJ; ISSN: 0021-8820, June 1997, XP002078543</p> <p>see page 530, right-hand column, paragraph 1 - page 531, left-hand column, paragraph 1; figure 3</p> <p style="text-align: center;">--- -/--</p>	1, 25, 27-32

☒ Further documents are listed in the continuation of box C.

☐ Patent family members are listed in annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

24 September 1998

Date of mailing of the international search report

08/10/1998

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Fuhr, C

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 98/09471

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>TSUGE K ET AL: "Characterization of Bacillus subtilis YB8, coproducer of lipopeptides surfactin and plipastatin B1."</p> <p>JOURNAL OF GENERAL AND APPLIED MICROBIOLOGY 41 (6). 1996. 541-545. ISSN: 0022-1260, XP002078544</p> <p>see page 541, paragraph 1</p> <p>see page 542, paragraph 1</p> <p>see page 542, last paragraph - page 543, paragraph 1</p> <p style="text-align: center;">---</p>	1,28-32
X	<p>YAMADA S ET AL: "BIOLOGICAL ACTIVITY OF ANTIFUNGAL SUBSTANCES PRODUCED BY BACILLUS-SUBTILIS."</p> <p>J PESTIC SCI 15 (1). 1990. 95-96. CODEN: NNGADV ISSN: 0385-1559, XP002078545</p> <p>see the whole document</p> <p style="text-align: center;">-----</p>	1,28-32